

Poster presentation

## Axis I-Axis II comorbidity of Obsessive Compulsive Personality Disorder: gender-related differences

Olga Georgiadou\*<sup>1</sup>, George Garyfallos<sup>2</sup>, Vasilis Bozikas<sup>2</sup>, Chrysi Protogerou<sup>1</sup>, Fotini Kosti<sup>1</sup>, Despina Ziliaskopoulou<sup>1</sup> and Aravella Adamopoulou<sup>1</sup>

Address: <sup>1</sup>Community Mental Health Centre, North-western District, Psychiatric Hospital of Thessaloniki, Greece and <sup>2</sup>2nd Department of Psychiatry, Aristotle University of Thessaloniki, Greece

\* Corresponding author

from International Society on Brain and Behaviour: 3rd International Congress on Brain and Behaviour Thessaloniki, Greece. 28 November – 2 December 2007

Published: 17 April 2008

*Annals of General Psychiatry* 2008, **7**(Suppl 1):S227 doi:10.1186/1744-859X-7-S1-S227

This abstract is available from: <http://www.annals-general-psychiatry.com/content/7/S1/S227>

© 2008 Georgiadou et al.; licensee BioMed Central Ltd.

### Background

Various studies validate that Obsessive Compulsive Personality Disorder is more common in men than in women. Most research concerning relations between criteria-defined OCPD and other disorders present significant comorbidity among them; which often inducts important diagnostic and treatment implications.

### Materials and methods

82 patients, 22 males and 60 females, who suffered from OCPD and have been treated at the Community Mental Health Centre within the past three years, were investigated. According to DSM-IV-TR criteria, patterns of Axis I and Axis II psychopathology were examined. Gender, age, education and family condition were also studied.

### Results

The proportion between men and women was almost 1:3. It is not surprising that patients were likely to have multiple diagnoses, 19 (86.4%) males and all females comorbid one or more Axis I disorders, also 13 (59.1%) males and 42 (70%) females comorbid one or more Axis II disorders. In particular 9 (40.9%) males and 35 (58.3%) females had mood disorders, with tendency for depression mostly for females (48.3 vs 27.3%) and tendency for dysthymic disorder mostly for males (13,6 vs 8.3%). High rates of comorbidity, 10 (45.5%) males and 32 (53.3%) females, have been also reported for anxiety disorders, with tendency for panic disorder with or without agora-

phobia mostly for females (35 vs 22.7%) and tendency for generalized anxiety disorder mostly for males (13,6 vs 6,7%). Somatoform and eating disorders frequently detected in females, in reverse impulse control disorders frequently detected in males. The findings support significant Cluster C, 10 (45.5%) males and 41 (68.3%) females, and Cluster B, 8 (36.4%) males and 25 (41.7%) females, personality disorders comorbidity. Dependent, histrionic, avoidant and borderline were mostly prevalent in females and respectively narcissistic in males. In addition, Cluster A personality disorders rarely co-occurred in both gender.

### Conclusions

The results from this study, suggest that the prevalence of DSM-IV-TR Axis I and Axis II disorders in our sample of OCPD patients and their gender-related differences, represent the Greek population who seek psychiatric support at the Community Mental Health Centre.

### References

1. Hopwood CJ, Morey LC, Gunderson JG, Skodol AE, Tracie Shea M, Grilo CM, McGlashan TH: **Hierarchical relationships between borderline, schizotypal, avoidant and obsessive-compulsive personality disorders.** *Acta Psychiatr Scand* 2006, **113**(5):430-439.
2. McGlashan TH, Grilo CM, Sanislow CA, Ralevski E, Morey LC, Gunderson JG, Skodol AE, Shea MT, Zanarini MC, Bender D, et al.: **Two-year prevalence and stability of individual DSM-IV criteria for schizotypal, borderline, avoidant, and obsessive-compulsive personality disorders: toward a hybrid model of axis II disorders.** *Am J Psychiatry* 2005, **162**(5):883-889.
3. Rossi A, Marnangeli MG, Butti B, Kalyvoka A, Petrucci C: **Pattern of comorbidity among anxious and odd personality disorders:**

the case of obsessive-compulsive personality disorder. *CNS Spectr* 2000, **5(9)**:23-26.

Publish with **BioMed Central** and every scientist can read your work free of charge

*"BioMed Central will be the most significant development for disseminating the results of biomedical research in our lifetime."*

Sir Paul Nurse, Cancer Research UK

Your research papers will be:

- available free of charge to the entire biomedical community
- peer reviewed and published immediately upon acceptance
- cited in PubMed and archived on PubMed Central
- yours — you keep the copyright

Submit your manuscript here:  
[http://www.biomedcentral.com/info/publishing\\_adv.asp](http://www.biomedcentral.com/info/publishing_adv.asp)

